

# Factorization of Symmetric Matrices and Actions on $\Sigma_k$

Baohua FU

February 8, 2008

## Abstract

In this note, firstly we give an easy proof of the factorization of symmetric matrices (see [Mos]), and then use it to prove the well-known fact that the automorphism group of a non-degenerate symmetric bilinear form  $Q$  acts transitively on the locus of isotropic subspaces  $\Sigma_k(Q)$ .

Let  $V$  be an  $m$ -dimensional complex vector space.

**Proposition 1.** *Every symmetric matrix  $Q \in \text{End}(V)$  is of the form  $P^T P$ , for some  $P \in \text{End}(V)$ .*

*Proof.* We will not make any difference between symmetric matrices and symmetric bilinear forms on  $V$ . If  $Q \neq 0$ , then there exists some  $u_1 \in V$  s.t.  $Q(u_1, u_1) \neq 0$ . Consider the hyperplane  $H_1 = \{u \in V | Q(u_1, u) = 0\}$ . If  $Q$  is not identically zero on  $H_1$ , then we can chose some  $u_2 \in H_1$ , s.t.  $Q(u_2, u_2) \neq 0$ , and then  $Q(u_2, u_1) = 0$ . Then we consider  $H_2 = \{u \in H_1 | Q(u_2, u) = 0\}$ , and we continue this procedure. At the end, we will get some vectors  $v_1, \dots, v_l, v_{l+1}, \dots, v_m$ , such that

$$Q(v_i, v_j) = \delta_{ij}, \text{ for } 1 \leq i, j \leq l; \quad Q(v_i, v_j) = 0, \text{ if } \max\{i, j\} \geq l + 1.$$

In this basis,  $Q$  is of the form  $\text{diag}(I_l, 0)$ , i.e. there exists some invertible matrix  $A$  such that  $A^T Q A = \text{diag}(I_l, 0)$ . Now we just take  $P = \text{diag}(I_l, 0) A^{-1}$ , then  $Q = P^T P$ .  $\square$

**Remark 1.** *This proposition is the principal result in [Mos] (theorem 2), proved by a much lengthier argument. Our proof works also for symmetric matrices over an algebraically closed field.*

Now let  $Q$  be a non-degenerate symmetric bilinear form on  $V$ . Recall that a  $k$ -plane  $\Lambda \subset V$  is said *isotropic* if  $Q|_{\Lambda \times \Lambda} = 0$ . Let  $\Sigma_k$  be the locus of isotropic subspaces of  $Q$ . It is well-known that  $\Sigma_k$  is not empty iff  $k \leq m/2$ , and irreducible if  $k < m/2$ . Let

$$G = \{A \in GL(V) \mid Q(Av, Aw) = Q(v, w) \text{ for all } v, w \in V\},$$

which acts naturally on  $\Sigma_k$ . The main purpose is to prove the following well-known fact (see [ACGH], p102):

**Proposition 2.** *The action of  $G$  on  $\Sigma_k$  is transitive.*

*Proof.* As easily seen, if  $k = m/2$ , then this proposition is true. In the following, we suppose  $k < m/2$ , thus  $\Sigma_k$  is irreducible. Since  $Q$  is non-degenerate, our previous proposition shows that we can suppose  $Q = I_m$ . Now  $G = \{A \in GL(V) \mid A^T A = I\}$ . Any  $k$ -plane is represented by an  $m \times k$  matrix  $\Lambda$ . We can suppose

$$\Lambda = \begin{pmatrix} I_k \\ M_1 \\ M_2 \end{pmatrix}$$

where  $M_1$  is a  $k \times k$  matrix, and  $M_2$  is an  $(m - 2k) \times k$  matrix. The isotropic condition is

$$I_k + M_1^T M_1 + M_2^T M_2 = 0.$$

By this we see that  $rk(M_1^T, M_2^T) = k$ . Using some matrix of the form  $I_m + E_{ij} + E_{ji} - E_{ii} - E_{jj} \in G$ , where  $E_{ij}$  is the matrix with 1 at the position  $(i, j)$  and 0 elsewhere, we can suppose  $M_1$  is invertible.

**Key claim:** For any  $\Lambda \in \Sigma_k$ , the orbit  $G \cdot \Lambda$  is dense in  $\Sigma_k$ .

*Proof of the key claim.* Take a matrix

$$A = \begin{pmatrix} I_k & 0 & 0 \\ 0 & A_1 & A_2 \\ 0 & B_1 & B_2 \end{pmatrix}$$

we will find out some conditions for  $\Lambda'$  such that there exists some  $A \in G$  of the above form with  $A\Lambda = \Lambda'$ . The last condition gives

$$A_1 = N_1 - A_2 M, \quad B_1 = N_2 - B_2 M$$

where  $N_1 = M'_1 M_1^{-1}$ ,  $N_2 = M'_2 M_1^{-1}$ ,  $M = M_2 M_1^{-1}$ . The isotropic conditions for  $\Lambda, \Lambda'$  imply that

$$N_1^T N_1 + N_2^T N_2 = I_k + M^T M.$$

The condition  $A^T A = I_m$  gives three equations. Combining with the above equation, we see that among the three there are only two independent ones, which can be simplified as

$$\begin{cases} N_1^T A_2 + N_2^T B_2 = M^T \\ A_2^T A_2 + B_2^T B_2 = I_{m-2k}. \end{cases}$$

From the first equation, we can resolve for  $A_2$ , and the only equation for  $B_2$  reads

$$B_2^T X B_2 - Y^T B_2 - B_2^T Y + Z = I_{m-2k},$$

where  $X, Z$  are symmetric,  $X = I_{m-2k} + N_2 N_1^{-1} (N_1^T)^{-1} N_2^T$ , so the condition  $rk(X) = m - 2k$  (maximal possible rank) is an open condition on  $\Lambda' \in \Sigma_k$ . If this condition is satisfied, our previous proposition gives  $X = P^T P$  for some invertible matrix  $P$ . Now the equation for  $B_2$  is equivalent to

$$(B_2^T P^T - Y^T P^{-1})(P B_2 - (P^{-1})^T Y) = W$$

where  $W$  is symmetric, thus of the form  $V^T V$ . Now we just take  $P B_2 - (P^{-1})^T Y = V$ , i.e.  $B_2 = P^{-1} V + X^{-1} Y$ . So if  $\Lambda'$  satisfies

$$rk(I_{m-2k} + M'_2 M_1'^{-1} (M_1'^T)^{-1} M_2'^T) = m - 2k,$$

then  $\Lambda' \in G \cdot \Lambda$ . As  $\Sigma_k$  is irreducible, we conclude that  $G \cdot \Lambda$  is dense in  $\Sigma_k$ .  $\square$

To finish the proof of the proposition, we just note that there exists at least one closed orbit, say  $G \cdot \Lambda$ . Then our key claim implies that  $G \cdot \Lambda = \Sigma_k$ , thus the action of  $G$  on  $\Sigma_k$  is transitive.  $\square$

*Acknowledgment:* This proof is done during the workshop on Algebraic Curves at UNSA, Nice. The author wants to thank J. Kock for reading a first draft of this note.

## References

- [Mos] A.Mostafazadeh *On a Factorization of Symmetric Matrices and Anti-linear Symmetries*, preprint math-ph/0203023.
- [ACGH] E.Arbarello, M.Cornalba, P.Griffiths, J.Harris, *Geometry of Algebraic Curves*, Springer-Verlag, 1985.

Labortoire J.A.Dieudonné, Parc Valrose  
06108 Nice cedex 02, FRANCE  
baohua.fu@polytechnique.org